


Examples

5.1 In the switching network shown, the switches operate independently. Each switch closes with probability p , and remains open with probability $1 - p$.

- a- Find the probability that a signal at the input will be rec
- b- Find the conditional probability that switch E is open, g



5.2 In a certain Village 20% of the population has disease D . A test is administered which has the property that if a person has D , the test will be positive 90% of the time, and if he does not have D , the test will be positive 30% of the time. All those whose test is positive are given a drug which invariably cures the disease, but produces a characteristic rash 25% of the time. Given that a person picked at random has the rash, what is the probability that he actually had D to begin with?

(Cumulative) Distribution Function

I. Discrete R. V.

Discrete R. V. X . pmf: $p_X(x_i) = P(\{X=x_i\})$

Define CDF $F_X(x) = P(\{X \leq x\}) = \sum_{x_i \leq x} p_X(x_i)$

Properties of $F_X(x)$:

- 1- $\lim_{x \rightarrow \infty} F_X(x) = 1$
- 2- $\lim_{x \rightarrow -\infty} F_X(x) = 0$
- 3- $P(\{a < X \leq b\}) = F_X(b) - F_X(a)$, i.e. $F_X(x)$ is a nondecreasing function.

II. Continuous R. V.

Define $F_X(x) = P(\{X \leq x\})$, and $f_X(x) = \frac{d}{dx}F_X(x) = \text{Probability Density Function.}$

$$P(\{a < X \leq b\}) = F_X(b) - F_X(a) = \int_a^b f_X(x) dx.$$

Properties of $f_X(x)$:

- 1- $f_X(x) \geq 0$
- 2- $\int_{-\infty}^{\infty} f_X(x) dx = 1$

Bernoulli Random Variable:

Experiment: "Bernoulli Trial" $S = \{\text{Success, Failure}\} = \{s, f\}$. $P(\{s\}) = p$, $P(\{f\}) = 1 - p$.

Bernoulli R.V. $X(\{\text{Success}\}) = 1$, $X(\{\text{Failure}\}) = 0$. $p_X(1) = p$, $p_X(0) = 1 - p$.

Binomial R.V. (Repeated Trials):

Experiment: N independent Bernoulli trials. $S = \{(s, s, \dots, s), (s, s, \dots, s, f), \dots, (f, f, \dots, f)\}$. 2^N sample points.

Define the **Binomial R. V.** by mapping each sample point into an integer (subset of reals) equal to the number of successes. How many points are there with n successes and N-n failures? $N!/n!(N-n)!$ Therefore

$$p_X(n) = \frac{N!}{(N-n)!n!} p^n (1-p)^{N-n} = \binom{N}{n} p^n (1-p)^{N-n} \text{ , where } \binom{N}{n} \equiv C(N, n) = \frac{N!}{(N-n)!n!} .$$

$$\text{Binomial Theorem: } (a + b)^N = \sum_{n=0}^N \binom{N}{n} a^n b^{N-n} .$$

$$\text{Binomial R.V. } X: p_X(n) = \binom{N}{n} p^n (1-p)^{N-n}, n = 0, 1, \dots, N .$$

$$\text{Then } \sum_{n=0}^N p_X(n) = \sum_{n=0}^N \binom{N}{n} p^n (1-p)^{N-n} = (p + (1-p))^N = 1 .$$

$$\binom{N}{n} = \text{number of ways of having } n \text{ successes and } N - n \text{ failures.}$$

$$\text{Then } \sum_{n=0}^N \binom{N}{n} = \text{total number of sample points} = (1 + 1)^N = 2^N .$$

For Ex. 2, $\{X = 1\} = \{H\}$, $\{X = 5\} = \{T, T, T, T, H\}$, $\{X < 3\} = \{(H), (T, H)\}$.

$$p_X(1) = 1/2, p_X(2) = 1/4, \dots, p_X(N) = 1/2^N .$$

For Ex. 3, $\{X = 3\} = \{(H, H, H, T), (H, H, T, H), (H, T, H, H), (T, H, H, H)\}$, $\{X = 0\} = \{T, T, T, T\}$.

$p_X(0) = 1/16$, $p_X(1) = 1/4$, $p_X(2) = 6/16 = 3/8$, $p_X(3) = 1/4$, $p_X(4) = 1/16$. For $p_X(2)$, the number of sample points with two heads is $4 \times 3/2! = 6$, we have total of 2^4 sample points of equal probability, hence the $6/16$ answer.

For Ex. 4, $S = \{X = 2\} + \{X = 3\} + \dots + \{X = 12\}$, a partition. Then $P(S) = p_X(2) + p_X(3) + \dots + p_X(12) = 1$.

Definition: A random variable that has only discrete experimental values (finite or countably infinite) is called a **Discrete Random Variable**.

Probability Mass Function (Probability Distribution):

Let X be a discrete R.V., and x be an experimental value. Define $p_X(x) = P\{X=x\}$ as the **probability mass function**.

A general property of probability mass function (pmf): $\sum_{\text{all } x} p_X(x) = 1$.

Random Variables

Definition:

A random variable, X , is a real-valued function defined on a sample space S . It is a mapping from S into \mathfrak{R} .

Examples:

Ex. 1: Toss coin. $S = \{H, T\}$. Define X such that $X(H) = 1$, $X(T) = 0$.

Ex. 2: Toss coin until head comes up. $S = \{H, (T, H), (T, T, H), \dots, (T, T, \dots, T, H)\}$.

Define $X(H) = 1$, $X(T, H) = 2$, $X(T, T, H) = 3$, \dots .

Ex. 3: Toss a coin 4 times for each experiment: $S = \{(H, H, H, H), (H, H, H, T), \dots, (T, T, T, T)\}$. Define X to be number of heads in sample point, e.g. $X(H, H, T, H) = 3$.

Ex. 4: Toss two dice: $S = \{(n, m): 1 \leq n, m \leq 6\}$, define $X((n, m)) = n + m$.

Events Defined From Random Variables:

Let X be a R.V. on S , $S = \{s_1, s_2, \dots\}$, defined by $X(s) = x$.

Let $\{X = 1\}$ define the event $\{s \in S: X(s) = 1\}$, similarly

let $\{X = a\} = \{s \in S: X(s) = a\}$, and

$\{a \leq X < b\} = \{s \in S: a \leq X(s) < b\}$.

For Ex. 1, $\{X = 1\} = \{H\}$, therefore $p_X(1) = 1/2$. $\{X = 0\} = \{T\}$, therefore $p_X(0) = 1/2$. $\{X = 2\} = \emptyset$, $\{X = \sqrt{3}\} = \emptyset$, $\{0 \leq X < 1\} = S$.